ELSEVIER

Journal of Organometallic Chemistry 651 (2002) 159

Journal of Organometallic Chemistry

www.elsevier.com/locate/jorganchem

Author Index of Volume 651

Adams, R.D., 124
Akiyama, Y., 137
Amour Morkan, İ., 132
Anulewicz-Ostrowska, R., 72

Belli Dell'Amico, D., 52
Bianchini, G., 5
Bohra, R., 98
Burzlaff, N., 66

Cai, X., 149, 150
Calderazzo, F., 52
Captain, B., 124
Castel, A., 44
Caubet, A., 105
Chavez, I., 44
Cingolani, A., 5

Delpech, F., 44
Drake, J.E., 98
Drozdov, A., 5

Erker, G., 157

Fischer, H., 66
Font-Bardía, M., 105
Fu, W., 124

Gamasa, M.P., 22
Gantzel, P., 90
Gehrhus, B., 149, 150
Gimeno, J., 22
Gornitzka, H., 44

Harada, A., 114
He, M., 146
Hitchcock, P.B., 149, 150
Hursthouse, M.B., 98

Isea, R., 22

Jain, V.K., 98
Joudat, M., 44

Kautz, J.A., 34
Kissounko, N.S., 34
Kissounko, D.A., 34
Kranenburg, M., 80
Kubiak, C.P., 90
Kwaśniewska, S., 72

Labella, L., 52
Lappert, M.F., 149, 150
Light, M.E., 98
López, C., 105

Manning, A.R., 60
Manriquez, J.M., 44
Marchetti, F., 5, 52
Martín-Vaca, B.M., 22
Meier, R.J., 80
Meijers, R.H.A.M., 80

Nakayama, Y., 114

Palmer, A.J., 60
Pérez, S., 105
Pettinari, C., 5

Ratnani, R., 98
Rivière, P., 44
Rouzaud, J., 44

Sbrana, E., 52
Sharma, R.K., 98
Sharma, V., 98
Simón-Manso, E., 90
Slootweg, J.C., 149, 150
Smith, M.D., 124
Solans, X., 105
Stone, F.G.A., 34
Stumpf, R., 66

Takashima, Y., 114
Takeuchi, R., 137
Tang, J., 146
Troyanov, S., 5

Vagedes, D., 157
Valderrama, M., 90
van der Pol, A., 80
van Heel, J.P.C., 80
Vegas, A., 22
Verkade, J.G., 15
Vertlib, V., 5

Weibert, B., 66
Wróblewski, R., 72

Yang, F., 146
Yang, J., 15
Yasuda, H., 114

Zhang, Y., 146
Zheng, R., 146

PII: S0022-328X(02)01453-5

ELSEVIER

Journal of Organometallic Chemistry 651 (2002) 161–163

Journal of Organometallic Chemistry

www.elsevier.com/locate/jorganchem

Subject Index of Volume 651

Activation
Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80
4-Acylpyrazolones
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]$Q^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5
Addition reactions
Non-catalyzed addition reactions of $Cl_3SiSiCl_3$ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15
Alkyls
The stable silylene Si[$(NCH^tBu)_2C_6H_4$-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{$Si(SiMe_3)_3$}$(THF)_3$] [R = Me, ^{t}Bu or $CH(SiMe_3)_2$], 150
Allylic phosphates
Iridium complex-catalyzed carbonylation of allylic phosphates, 137
π-Allyl iridium
Iridium complex-catalyzed carbonylation of allylic phosphates, 137
Aluminium
Steric effects in alkylalane dioldiates: [^{t}BuAl($OC(CH_3)_2CH_2C(CH_3)_2O$)]$_2$ a new (monoalkyl)alane *O*,*O*′-chelate complex, 72
Amino-functionalised silyl anions
The stable silylene Si[$(NCH^tBu)_2C_6H_4$-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{$Si(SiMe_3)_3$}$(THF)_3$] [R = Me, ^{t}Bu or $CH(SiMe_3)_2$], 150
Amino-substituents
Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe_2-2-Cl-2-PPh_3-*closo*-2,1-$RhCB_{10}H_{10}$] and [1-NH_3-2-Cl-2-PPh_3-*closo*-2,1-$RhCB_{10}H_{10}$], 34

Benzyne
CH and PC bond activations of PMe_2Ph ligands by an octahedral Ru_6 cluster, 124

Carbene complexes
The first carbene-*C*,*N* chelate tetracarbonyl dihalogeno and carbene-*C*,*O* chelate tetracarbonyl benzochinone tungsten complexes, 66
Carbonyl
High order cycloaddition reactions of $M(CO)_3$-coordinated *N*-cyanoazepine with alkynes; M: Cr, Mo, W, 132
Carbonylation
Iridium complex-catalyzed carbonylation of allylic phosphates, 137
Carborane
Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe_2-2-Cl-2-PPh_3-*closo*-2,1-$RhCB_{10}H_{10}$] and [1-NH_3-2-Cl-2-PPh_3-*closo*-2,1-$RhCB_{10}H_{10}$], 34
Catalytic yield
Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80
C–H activation
Heterodimetallic palladium(II) compounds containing terdentate [Csp^2(ferrocene),N,N′]$^-$ ferrocenyl ligands, 105
Chalcogen-bridged bis(aryloxo) ligands
Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114
Chalcogenide compounds
Chiral sulfur diphosphazanes derived from S-$(Ph_2P)_2N$(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of $Ph_2P(S)N(CHMePh)PPh_2$, {$Ph_2P(S)$}$_2$N-(CHMePh) and [(Cp*)MCl{η^2-P,S-$Ph_2PNHP(S)Ph_2$}]BF_4, Cp* = η^5-C_5Me_5; M = Rh, Ir, 90
Chelate complexes
The first carbene-*C*,*N* chelate tetracarbonyl dihalogeno and carbene-*C*,*O* chelate tetracarbonyl benzochinone tungsten complexes, 66
Chiral diphosphazanes
Chiral sulfur diphosphazanes derived from S-$(Ph_2P)_2N$(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of $Ph_2P(S)N(CHMePh)PPh_2$, {$Ph_2P(S)$}$_2$N-(CHMePh) and [(Cp*)MCl{η^2-P,S-$Ph_2PNHP(S)Ph_2$}]BF_4, Cp* = η^5-C_5Me_5; M = Rh, Ir, 90
Chromium
High order cycloaddition reactions of $M(CO)_3$-coordinated *N*-cyanoazepine with alkynes; M: Cr, Mo, W, 132
Cluster
CH and PC bond activations of PMe_2Ph ligands by an octahedral Ru_6 cluster, 124
Substitution reactions of [{$Co(\eta^5$-$C_5H_5)$}$_2${$Fe(CO)_2(PPh_3)$}(μ_3-S)(μ_3-CS)], 60
Cobalt
Substitution reactions of [{$Co(\eta^5$-$C_5H_5)$}$_2${$Fe(CO)_2(PPh_3)$}(μ_3-S)(μ_3-CS)], 60
Crystal structures
Chiral sulfur diphosphazanes derived from S-$(Ph_2P)_2N$(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of $Ph_2P(S)N(CHMePh)PPh_2$, {$Ph_2P(S)$}$_2$N-(CHMePh) and [(Cp*)MCl{η^2-P,S-$Ph_2PNHP(S)Ph_2$}]BF_4, Cp* = η^5-C_5Me_5; M = Rh, Ir, 90
Cycloaddition
New linked di-germanocenes and di-stannocenes, 44
Cyclometallation
CH and PC bond activations of PMe_2Ph ligands by an octahedral Ru_6 cluster, 124
Heterodimetallic palladium(II) compounds containing terdentate [Csp^2(ferrocene),N,N′]$^-$ ferrocenyl ligands, 105
Cyclopalladated complex
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146

1,2-Diketones
Non-catalyzed addition reactions of $Cl_3SiSiCl_3$ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15
β-Diketones
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of

PII: S0022-328X(02)01454-7

[Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q$^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5

Diols
Steric effects in alkylalane dioldiates: [^{t}BuAl(OC(CH$_3$)$_2$CH$_2$C-(CH$_3$)$_2$O)]$_2$ a new (monoalkyl)alane *O*,*O*′-chelate complex, 72

Disilane
Non-catalyzed addition reactions of $Cl_3SiSiCl_3$ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15

Ferrocenylimines
Heterodimetallic palladium(II) compounds containing terdentate [Csp^2(ferrocene),N,N′]$^-$ ferrocenyl ligands, 105

Functionalized oxime
Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [{R$_2$Sn(ON=C(Me)py)}$_2$O]$_2$ (R = Bu^n and Et) and 2-NC$_5$H$_4$C(Me)=NOH, 98

Germanate
New linked di-germanocenes and di-stannocenes, 44

Germanocene
New linked di-germanocenes and di-stannocenes, 44

Group 6 transition metal complexes
Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114

Heck reaction
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146

Indenyl complexes
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=$\overline{COCH_2(CH_2)_3C}H_2$}($\eta^5$-C$_9H_7$)(PPh$_3$)$_2$][PF$_6$]·CH$_2Cl_2$, 22

Iridium
Iridium complex-catalyzed carbonylation of allylic phosphates, 137

Iron
Substitution reactions of [{Co(η^5-C$_5$H$_5$)}$_2${Fe(CO)$_2$(PPh$_3$)}(μ_3-S)(μ_3-CS)], 60

Ligand substitution
Substitution reactions of [{Co(η^5-C$_5$H$_5$)}$_2${Fe(CO)$_2$(PPh$_3$)}(μ_3-S)(μ_3-CS)], 60

Lithium
The stable silylene Si[(NCH$_2^t$Bu)$_2$C$_6$H$_4$-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{Si(SiMe$_3$)$_3$}(THF)$_3$] [R = Me, ^{t}Bu or CH(SiMe$_3$)$_2$], 150

Methylation
Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80

Multinuclear NMR
Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [{R$_2$Sn(ON=C(Me)py)}$_2$O]$_2$ (R = Bu^n and Et) and 2-NC$_5$H$_4$C(Me)=NOH, 98

N-Cyanoazepine
High order cycloaddition reactions of M(CO)$_3$-coordinated *N*-cyanoazepine with alkynes; M: Cr, Mo, W, 132

N,*N*-dialkylcarbamato
Organometallic complexes of ruthenium(II) with *N*,*N*-dialkylcarbamato as the supporting ligand, 52

N,N′Ferrocenyl ligands
Heterodimetallic palladium(II) compounds containing terdentate [Csp^2(ferrocene),N,N′]$^-$ ferrocenyl ligands, 105

Organometallic complexes
Organometallic complexes of ruthenium(II) with *N*,*N*-dialkylcarbamato as the supporting ligand, 52

Ortho-metallation
CH and PC bond activations of PMe$_2$Ph ligands by an octahedral Ru$_6$ cluster, 124

Oxacycloalkenyl complexes
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=$\overline{COCH_2(CH_2)_3C}H_2$}($\eta^5$-C$_9H_7$)(PPh$_3$)$_2$][PF$_6$]·CH$_2Cl_2$, 22

Oxacycloalkylidene complexes
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=$\overline{COCH_2(CH_2)_3C}H_2$}($\eta^5$-C$_9H_7$)(PPh$_3$)$_2$][PF$_6$]·CH$_2Cl_2$, 22

Oxidation
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q$^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5

Oxidative decarbonylation
The first carbene-*C*,*N* chelate tetracarbonyl dihalogeno and carbene-*C*,*O* chelate tetracarbonyl benzochinone tungsten complexes, 66

Palladium catalyst
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146

Palladium mirror
Cyclopalladated complexes of tertiary arylamines as highly efficient catalysts using in the Heck reactions, 146

P–C cleavage
CH and PC bond activations of PMe$_2$Ph ligands by an octahedral Ru$_6$ cluster, 124

Phosphines
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q$^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5

^{31}P-NMR
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]Q$^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5

1,4-Quinone
Non-catalyzed addition reactions of $Cl_3SiSiCl_3$ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15

1,2-Quinones
Non-catalyzed addition reactions of $Cl_3SiSiCl_3$ with 1,2-diketones, 1,2-quinones and with a 1,4-quinone, 15

Rhodacarboranes
Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe$_2$-2-Cl-2-PPh$_3$-*closo*-2,1-RhCB$_{10}$H$_{10}$] and [1-NH$_3$-2-Cl-2-PPh$_3$-*closo*-2,1-RhCB$_{10}$H$_{10}$], 34

Rhodium
Synthesis and reactivity of the rhodium monocarbollide complexes [1-NH=CMe$_2$-2-Cl-2-PPh$_3$-*closo*-2,1-RhCB$_{10}$H$_{10}$] and [1-NH$_3$-

2-Cl-2-PPh_3-*closo*-2,1-$RhCB_{10}H_{10}$], 34
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]$Q^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5
Rhodium and Iridium complexes
Chiral sulfur diphosphazanes derived from S-$(Ph_2P)_2N$(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of $Ph_2P(S)N(CHMePh)PPh_2$, $\{Ph_2P(S)\}_2N$-(CHMePh) and [(Cp*)MCl{η^2-P,S-$Ph_2PNHP(S)Ph_2$}]BF_4, Cp* = η^5-C_5Me_5; M = Rh, Ir, 90
Ring-opening metathesis polymerization
Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114
Ruthenium
CH and PC bond activations of PMe_2Ph ligands by an octahedral Ru_6 cluster, 124
Synthesis and characterization of 2-oxacycloalkylideneruthenium(II) complexes: X-ray crystal structure of the 2-oxacycloheptylidene complex [Ru{=$\overline{C}OCH_2(CH_2)_3\overline{C}H_2$}($\eta^5$-$C_9H_7$)$(PPh_3)_2$][$PF_6$]·$CH_2Cl_2$, 22
Ruthenium(II)
Organometallic complexes of ruthenium(II) with *N*,*N*-dialkylcarbamato as the supporting ligand, 52

SET reaction
New linked di-germanocenes and di-stannocenes, 44
Sisyls
The stable silylene Si[$(NCH_2^tBu)_2C_6H_4$-1,2]: insertion into Li–C or Li–Si bonds of lithium alkyls LiR or [Li{$Si(SiMe_3)_3$}$(THF)_3$] [R = Me, ^{t}Bu or $CH(SiMe_3)_2$], 150
Stannocene
New linked di-germanocenes and di-stannocenes, 44
Stereo selectivity
Synthesis of novel oxo complexes of tungsten and molybdenum with various chalcogen-bridged chelating bis(aryloxo) ligands and their catalytic behavior for ring-opening metathesis polymerization, 114
Steric effects
Steric effects in alkylalane dioldiates: [^{t}BuAl($OC(CH_3)_2CH_2$-$C(CH_3)_2O$)]$_2$ a new (monoalkyl)alane *O*,*O*′-chelate complex, 72
Structures
The reactivity of new (1,5-cyclooctadiene)rhodium acylpyrazolonates towards N- and P-donor ligands: X-ray structures of [Rh(1,5-COD)Q^s], [Rh(1,5-COD)(phen)]$Q^s \cdot 0.5H_2O$ (HQ^s = 1-phenyl-3-methyl-4-(2-thenoyl)-pyrazol-5-one) and [Rh(1,5-COD)Br]$_2$, 5
syn–*anti* Isomerization
Iridium complex-catalyzed carbonylation of allylic phosphates, 137
Synthesis
Chiral sulfur diphosphazanes derived from S-$(Ph_2P)_2N$(CHMePh) and its rhodium(I), (III) and iridium(III) complexes. Crystal structures of $Ph_2P(S)N(CHMePh)PPh_2$, $\{Ph_2P(S)\}_2N$-(CHMePh) and [(Cp*)MCl{η^2-P,S-$Ph_2PNHP(S)Ph_2$}]BF_4, Cp* = η^5-C_5Me_5; M = Rh, Ir, 90

Tetraorganodistannoxane
Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [{$R_2Sn(ON{=}C(Me)py)\}_2O]_2$ (R = Bu^n and Et) and 2-$NC_5H_4C(Me){=}NOH$, 98
Thiocarbonyl
Substitution reactions of [{$Co(\eta^5$-$C_5H_5)\}_2${$Fe(CO)_2(PPh_3)$}(μ_3-S)(μ_3-CS)], 60
Tungsten complexes
The first carbene-*C*,*N* chelate tetracarbonyl dihalogeno and carbene-*C*,*O* chelate tetracarbonyl benzochinone tungsten complexes, 66

β,γ-Unsaturated esters
Iridium complex-catalyzed carbonylation of allylic phosphates, 137

Vibrational spectroscopy
Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80

X-ray structures
Synthesis, spectroscopic and structural aspects of some tetraorganodistannoxanes with internally functionalized oxime: crystal and molecular structures of [{$R_2Sn(ON{=}C(Me)py)\}_2O]_2$ (R = Bu^n and Et) and 2-$NC_5H_4C(Me){=}NOH$, 98
Zirconocene
Characterization of a metallocene/co-catalyst system supported on silica by Fourier-Transform Raman spectroscopy, 80